

Lec 02: Sets II

Prof. Adam J. Aviv

GW

CSCI 1311 Discrete Structures I
Spring 2023

Sets, so far...

- Sets are a collection of unique items, including other sets, that are designated using braces.
 - ▶ $\{1, 2, 3\}$, $\{\text{red}, \text{blue}\}$, $\{\}$, $\{5, \{4, 3\}, 8\}$
- We described an element as a member of a set using \in or not as a member using \notin .
 - ▶ $1 \in \{1, 2, 3\}$ and $7 \notin \{1, 2, 3\}$
- Two sets are equal = if they contain the same elements, the order is irrelevant.
 - ▶ $\{1, 2, 3\} = \{2, 1, 3\}$ and $\{1, 2, 3\} \neq \{2, 3, 4\}$
- Set builder notation: $\{x \in S \mid P(x)\}$
 - ▶ $\{x \in \mathbb{Z} \mid x > 0 \text{ and } x \neq 15\}$
- Cardinality of a set is the number of elements contained in the set
 - ▶ $|\{1, 2, 3\}| = 3$ $|\{1, \{2, 3\}\}| = 2$ $|\{x \in \mathbb{Z} \mid x > -5\}| = \infty$
- If a the cardinality of a set is zero, we call that the empty set
 - ▶ $|\{\}| = 0$, $|\emptyset| = 0$
 - ▶ But the empty set can be member of a set, $|\{\{\}\}| = |\{\emptyset\}| = 1$

Today

- Subsets
- Union, Intersection, Complement and Difference
- Partitions
- Power Sets
- Order Pairs and Tuples
- Cartesian Products

Subsets

Definition

For two sets A and B , A is a **subset** of B (written $A \subseteq B$) if for all elements $x \in A$ then $x \in B$.

Example

For the set $A = \{9, 7, 22\}$ and $B = \{9, 22, 18, 42, 7\}$, $A \subseteq B$ (or $B \supseteq A$) because all the elements of A are also in B , that is $9 \in B$, $7 \in B$ and $22 \in B$.

Definition

A **proper subset** (written $A \subset B$) is a subset whereby $A \subseteq B$ but there exists at least one element $x \in B$ such that $x \notin A$.

Exercises

For the set $A = \{42, 18, 3, 22\}$ and $B = \{22, 3, 42, 18\}$,

is $A \subset B$?

Is $A \subseteq B$?

For two sets, C and D , where $C = D$, is $C \subseteq D$?

For two sets E and F , if $E \subseteq F$ and $E \supseteq F$, then it must be the case that $E = F$. Why?

Is $\emptyset \subset \{5, 12, 2, 19\}$?

Is $\emptyset \subseteq \emptyset$?

Union and Intersection

Definition

The **union** of two sets A and B , written $A \cup B$, is the set that contains all elements in A *or* B .

Definition

The **intersection** of two sets A and B , written $A \cap B$, is the set that contains all elements in A *and* B .

Example

If $A = \{12, 7, 19, 5\}$ and $B = \{12, 3, 22, 42, 5\}$ then

$$A \cup B = \{12, 7, 19, 5, 3, 22, 42\} \text{ and}$$

$$A \cap B = \{12, 5\}$$

Complements or Difference

Definition

The **difference** between set B and A (or the relative complement of A in B), written $B \setminus A$ or $B - A$, is the set of elements in B that are not in A .

Example

If $A = \{12, 7, 19, 5\}$ and $B = \{12, 3, 22, 42, 5\}$ then


$$A \setminus B = A - B = \{7, 19, 5\} \text{ and}$$

$$B \setminus A = B - A = \{3, 22, 42\}$$

Definition

The **compliment** of set A , written A^c is the set of all elements in the universe U from which sets are populated that are not in A .

Set Operators as Venn Diagrams



Exercises

For any set A , what is

$$A \cup \emptyset?$$

$$A \cap \emptyset?$$

$$A \cup A?$$

$$A \cap A?$$

Write the set builder notation equivalent for each operation, assuming a universal set U .

$$A \cup B$$

$$A \cap B$$

$$B - A$$

$$A^c$$

Operators on Indexed Collections of Sets

Consider an *indexed* collection of sets $A_1, A_2, A_3, \dots, A_n$ where n is a positive integer, then we can define the following index operations over the sets:

- $\bigcup_{i=1}^n A_i = A_1 \cup A_2 \cup A_3 \cup \dots \cup A_n = \{x \mid x \in A_j \text{ for some } j = 1, 2, \dots, n\}$
- $\bigcap_{i=1}^n A_i = A_1 \cap A_2 \cap A_3 \cap \dots \cap A_n = \{x \mid x \in A_j \text{ for all } j = 1, 2, \dots, n\}$

We can even define these up to ∞ :

- $\bigcup_{i=1}^{\infty} A_i = A_1 \cup A_2 \cup A_3 \cup \dots = \{x \mid x \in A_j \text{ for some } j = 1, 2, \dots\}$
- $\bigcap_{i=1}^{\infty} A_i = A_1 \cap A_2 \cap A_3 \cap \dots = \{x \mid x \in A_j \text{ for all } j = 1, 2, \dots\}$

Mutually Disjoint Sets

Definition

Collection of indexed sets A_1, A_2, A_3, \dots are called **mutually disjoint** (or **pairwise disjoint**) if and only if for every two sets A_i and A_j , $A_i \cap A_j = \emptyset$

Example

The following sets are mutually disjoint

- $N_0 = \{n \mid n = 3k \text{ for some } k \in \mathbb{N}\} = \{0, 3, 6, \dots\}$
- $N_1 = \{n \mid n = 3k + 1 \text{ for some } k \in \mathbb{N}\} = \{1, 4, 7, \dots\}$
- $N_2 = \{n \mid n = 3k + 2 \text{ for some } k \in \mathbb{N}\} = \{2, 5, 8, \dots\}$

Partition

Definition

Let A_1, A_2, A_3, \dots be a **partition** of the set A if A_1, A_2, A_3, \dots are mutually disjoint and $\bigcup_i A_i = A$

Example

The following sets are a partition of \mathbb{N} (natural numbers)

- $N_0 = \{n \mid n = 3k \text{ for some } k \in \mathbb{N}\} = \{0, 3, 6, \dots\}$
- $N_1 = \{n \mid n = 3k + 1 \text{ for some } k \in \mathbb{N}\} = \{1, 4, 7, \dots\}$
- $N_2 = \{n \mid n = 3k + 2 \text{ for some } k \in \mathbb{N}\} = \{2, 5, 8, \dots\}$

Because N_0, N_1, N_2 are mutually disjoint and $N_0 \cup N_1 \cup N_2 = \mathbb{N}$

Exercises

Write down three sets that are mutually disjoint

Define a partition for \mathbb{Z}

Is the following a valid partition of the set $M = \{1, 2, 3, \dots, 9\}$:

- $\{x \in M \mid x \notin \{1, 2, 3\}\}$
- $\{x \in M \mid x \notin \{4, 5, 6\}\}$
- $\{x \in M \mid x \notin \{7, 8, 9\}\}$

Power Sets

Definition

The **power set** of A , written $\mathcal{P}(A)$, is the set of all subsets of A .

Example

The power set of $A = \{1, 2, 3\}$ is

$$\begin{aligned}\mathcal{P}(A) = \{ & \emptyset, \\ & \{1\}, \{1, 2\}, \{1, 3\}, \{1, 2, 3\}, \\ & \{2\}, \{2, 3\}, \\ & \{3\} \end{aligned}$$

The cardinality of a power set, $|\mathcal{P}(A)|$ is a power of 2, namely, $|\mathcal{P}(A)| = 2^{|A|}$

Why?

The empty set \emptyset is a member of any power set, why?

Because the size of the power set is always a power of 2, we also denote the power set of a set A as 2^A

Example

$$\mathcal{P}(\{a, b\}) = 2^{\{a,b\}} = \{\emptyset, \{a\}, \{b\}, \{a, b\}\}$$

Ordered Pairs

Definition

An **ordered pair** of two elements a and b , written (a, b) is a grouping of a and b where a is the *first element* and b is the *second element*.

Definition

Two ordered pairs are equal, written $(a, b) = (c, d)$, if and only if $a = b$ and $c = d$.

Example

An ordered pair is used to define the Cartesian plane, where the first element is the x component and the second element is the y component.

Two points in the plain are only equal when they have the same x and y

Generalizing ordered pairs into tuples

We can define ordered collections of more than two elements by combining ordered pairs. For example, an ordered collection of three elements and four elements can be written:

$$x = (a, (b, c)) \quad y = (a, (b, (c, d)))$$

As this is cumbersome, we can combine these elements together to form a **tuple**, denoted like so

$$x = (a, b, c) \quad y = (a, b, c, d)$$

A tuple has the same properties as an ordered pair, where each element of the two tuples must be equal for the tuples to be equal. For example,

$$(10, 12, 3, 5) \neq (10, 2, 3, 5)$$

Cartesian Products (or Cross Products)

Definition

The **Cartesian product** (or **cross product**) of two sets A and B , written $A \times B$ and read “A cross B”, is the set of all ordered pairs (a, b) where $a \in A$ and $b \in B$.

Example

We can denote the cross product of two sets using set builder notation:

$$A \times B = \{(a, b) \mid a \in A \text{ and } b \in B\}$$

We can further generalize the result for multiple products,

$$A_1 \times A_2 \times A_3 \times \dots = \{(a_1, a_2, a_3, \dots) \mid a_1 \in A_1, a_2 \in A_2, a_3 \in A_3, \dots\}$$

Exercises

Let $A = \{1, 2, 3\}$ and $B = \{u, v\}$, what is:

$A \times B$

$B \times A$

$A \times A$

$B \times B$

Let $D = \{p, q\}$, what is $2^D \times B$?